

KOMANEKA *at Bisma*

KODOK SAWAH

Rice Field Frog

Fejervarya cancrivora


Kodok Sawah is an amphibian that lives a lot in rice fields, swamps, and ditches to mangrove swamps. They have rough skin and nodules or knuckles, often dry, and their hind legs are often short, so that most frogs are not good at jumping far.

Adult males are about **60 mm** and mature females are around **70-80 mm** and weigh about **1.5 to 3 ounces**.

In general, they are harmless animals.

Their favorite time is rainy season, they will make sounds and shout back and forth with other frogs.

Latin Name : *Fejervarya cancrivora*



TOKEK

Gecko

Gekkonidae

Gekkonidae is a class/family of lizards which includes the types of geckos and lizards that are spread throughout the tropical regions of the world. Several types of this tribe include the most common types of lizards found and live around humans. They have a larger body size than lizards. The total length reaches **30 cm**, almost half of which is the length of the tail.

Gekkonidae or ***Tokek*** have a life span of between **10 – 20 years**.

They are not dangerous unless there is an outside threat.

Because geckos have a fairly high level of light sensitivity, they don't like bright places. That's why, they often appear or nest in the corners of a small room and lack of light.

Latin Name : *Gekkonidae*



KOMANEKA *at Bisma*

CICAK

House Gecko

Hemidactylus frenatus

The common **House Gecko** *Hemidactylus frenatus* is a gecko native to South and Southeast Asia. They grow to a length of between **7.5–15 cm (3–6 in)**, and live for about **7 years**. These small geckos are non-venomous and not harmful to humans.

The common house gecko is tropical, and thrives in warm, humid areas where it can crawl around on rotting wood in search of the insects it eats, as well as within urban landscapes in warm climates. In Asia/Southeast Asia, geckos have local names **onomatopoeically** derived from the sounds they make:

Hemidactylus frenatus is called “chee chak” or “chi chak” (pr- chee chuck), said quickly, also commonly spelled as “cicak” in Malay dictionaries. They will shed their tails when threatened by predators and can regenerate their tails in **1 month**.

Latin Name : *Hemidactylus frenatus*

